Miltenyi Biotec

ODN 2088 Control (ODN 2087)

Order no. 130-105-819
Order no. 130-105-940

Contents

1. Description
2. Background information
3. Applications
   - 3.1 General applications
   - 3.2 Specific applications
4. Instructions for use
   - 4.1 Recommended concentrations
   - 4.2 Reconstitution protocol
5. References

1. Description

Components	**130-105-819:** 200 µg lyophilized ODN 1 mL 1× TE Buffer or **130-105-940:** 1 mg lyophilized ODN 1 mL 1× TE Buffer.
Description	TLR7/8 antagonist, TLR9 antagonist control.
Product format	Lyophilized product without carrier protein or preservatives.
Sequence	dT*dC*dC*dT*dG*dA*dG*dC*dT*dT*dG*dA*dA*dG*dT * Phosphorothioate backbone
Endotoxin level	Low endotoxin (<1 EU/mg) as determined by kinetic Limulus Amebocyte Lysate (LAL) assay.
Storage	Store lyophilized product at –20 °C. Upon reconstitution, aliquots should be stored at –20 °C and are stable for 6 months. Avoid repeated freeze-thaw cycles. The expiration date is indicated on the vial label.

2. Background information

TLR7, TLR8, and TLR9 are prominent members of the Toll-like-receptor (TLR) family which recognize nucleic acids as pathogen-associated molecular patterns.[3] Activation of these TLRs can be inhibited by short single-stranded synthetical oligodeoxyribonucleotides (ODNs).[4,6] Several classes of and motifs for inhibitory ODN have been identified.[1,6,7]

3. Applications

3.1 General applications

- TLR7/8/9 antagonists can be used to inhibit TLR7/8/9-mediated cellular responses, such as activation of immune cells, human PBMCs, murine splenocytes, or isolated immune cells, e.g., B cells and pDCs.
- TLR7, TLR8, or TLR9 antagonists can be used to inhibit signaling in TLR7/8/9-expressing recombinant cell lines.

3.2 Specific applications

- ODN 2088 Control (ODN 2087) can be used as a sequence control for ODN 2088.[1,2,4-7] ODN 2088 Control (ODN 2087) inhibits TLR7- and TLR8-mediated signaling but not TLR9-mediated signaling.[5]

4. Instructions for use

4.1 Recommended concentrations

Recommended concentrations for use are

for human and murine immune cells: 0.05–5 µM

for recombinant cell lines: 0.05–10 µM

The optimal concentration range needs to be tested for individual assay systems.

4.2 Reconstitution protocol

1. Spin down pellet.
2. a) For 200 µg lyophilized ODN:
   To obtain a 200 µM solution resuspend pellet in 208 µL of 1× TE Buffer.
   ▲ **Note:** Alternatively, PBS or water can be used for reconstitution.

   b) For 1 mg lyophilized ODN:
   To obtain a 200 µM solution resuspend pellet in 1040 µL of 1× TE Buffer.
   ▲ **Note:** Alternatively, PBS or water can be used for reconstitution.
3. Vortex and incubate overnight at 4 °C.
4. Store aliquots at –20 °C.

Miltenyi Biotec B.V. & Co. KG
Friedrich-Ebert-Straße 68, 51429 Bergisch Gladbach, Germany
Phone +49 2204 8306-0, Fax +49 2204 85197
macsde@miltenyi.com
www.miltenyibiotec.com

140-004-795.02

5. References

1. Hackstein, H. *et al.* (2011) The TLR7/8 ligand resiquimod targets monocyte-derived dendritic cell differentiation via TLR8 and augments functional dendritic cell generation. Cell Immunol. 271: 401–412.
2. Jurk, M. *et al.* (2006) Modulating responsiveness of human TLR7 and 8 to small molecule ligands with T-rich phosphorothiate oligodeoxynucleotides. Eur. J. Immunol. 36: 1815–1826.
3. Kawai, T. and S. Akira (2010) The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. Nat. Immunol. 11: 373–384.
4. Krieg, A. M. *et al.* (1998) Sequence motifs in adenoviral DNA block immune activation by stimulatory CpG motifs. Proc. Natl. Acad. Sci. U. S. A. 95: 12631–12636.
5. Lenert, P. *et al.* (2001) CpG stimulation of primary mouse B cells is blocked by inhibitory oligodeoxyribonucleotides at a site proximal to NF-kappaB activation. Antisense Nucleic Acid Drug Dev. 11: 247–256.
6. Lenert, P. S. (2010) Classification, mechanisms of action, and therapeutic applications of inhibitory oligonucleotides for Toll-like receptors (TLR) 7 and 9. Mediators Inflamm. 2010: 986596.
7. Stunz, L. L. *et al.* (2002) Inhibitory oligonucleotides specifically block effects of stimulatory CpG oligonucleotides in B cells. Eur. J. Immunol. 32: 1212–1222.

Refer to **www.miltenyibiotec.com** for all data sheets and protocols. Miltenyi Biotec provides technical support worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

Legal notices

Limited product warranty

Miltenyi Biotec B.V. & Co. KG and/or its affiliate(s) warrant this product to be free from material defects in workmanship and materials and to conform substantially with Miltenyi Biotec's published specifications for the product at the time of order, under normal use and conditions in accordance with its applicable documentation, for a period beginning on the date of delivery of the product by Miltenyi Biotec or its authorized distributor and ending on the expiration date of the product's applicable shelf life stated on the product label, packaging or documentation (as applicable) or, in the absence thereof, ONE (1) YEAR from date of delivery ("Product Warranty"). Miltenyi Biotec's Product Warranty is provided subject to the warranty terms as set forth in Miltenyi Biotec's General Terms and Conditions for the Sale of Products and Services available on Miltenyi Biotec's website at www.miltenyibiotec.com, as in effect at the time of order ("Product Warranty"). Additional terms may apply. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING IF A PRODUCT IS SUITABLE FOR CUSTOMER'S PARTICULAR PURPOSE AND APPLICATION METHODS.

Technical information

The technical information, data, protocols, and other statements provided by Miltenyi Biotec in this document are based on information, tests, or experience which Miltenyi Biotec believes to be reliable, but the accuracy or completeness of such information is not guaranteed. Such technical information and data are intended for persons with knowledge and technical skills sufficient to assess and apply their own informed judgment to the information. Miltenyi Biotec shall not be liable for any technical or editorial errors or omissions contained herein.
All information and specifications are subject to change without prior notice. Please contact Miltenyi Biotec Technical Support or visit www.miltenyibiotec.com for the most up-to-date information on Miltenyi Biotec products.

Licenses

This product and/or its use may be covered by one or more pending or issued patents and/or may have certain limitations. Certain uses may be excluded by separate terms and conditions. Please contact your local Miltenyi Biotec representative or visit Miltenyi Biotec's website at www.miltenyibiotec.com for more information.
The purchase of this product conveys to the customer the non-transferable right to use the purchased amount of the product in research conducted by the customer (whether the customer is an academic or for-profit entity). This product may not be further sold. Additional terms and conditions (including the terms of a Limited Use Label License) may apply.
CUSTOMER'S USE OF THIS PRODUCT MAY REQUIRE ADDITIONAL LICENSES DEPENDING ON THE SPECIFIC APPLICATION. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING FOR ITSELF WHETHER IT HAS ALL APPROPRIATE LICENSES IN PLACE. Miltenyi Biotec provides no warranty that customer's use of this product does not and will not infringe intellectual property rights owned by a third party. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

Trademarks

The Miltenyi Biotec logo is a registered trademark or trademark of Miltenyi Biotec and/or its affiliates in various countries worldwide.

Copyright © 2020 Miltenyi Biotec and/or its affiliates. All rights reserved.