Miltenyi Biotec

Mouse IL-7-Fc
research grade

Contents

1. Description
   - 1.1 Background information
   - 1.2 Applications
2. References

1. Description

Products

Mouse IL-7-Fc, research grade.

Fc-fused recombinant mouse interleukin 7.

Content in µg	Order no.
10	130-137-043
25	130-137-018

Biological activity

The ED_{50} is ≤20 ng/mL corresponding to an activity of ≥5×10^4 U/mg.

▲ **Note:** The ED50 is determined by proliferation assay using mouse 2E8 cells.

Primary structure

Two identical glycosylated covalently-linked polypeptide chains (363 amino acid residues each).

Molecular mass

82.0 kDa (dimer, calculated).

Source

Produced in HEK293 cells.

Product format

Lyophilized from a filtered (0.2 µm) buffer solution.

Stabilizer

Mannitol and trehalose.

Purity

>95% as determined by SDS-PAGE analysis.

Endotoxin level

Low endotoxin (<1.0 EU/µg cytokine) as determined by Limulus Amebocyte Lysate (LAL) assay.

Storage

Lyophilized Mouse IL-7-Fc, research grade should be stored at –20 °C. The expiration date is indicated on the vial label. Upon reconstitution aliquots should be stored at –20 °C or below. Avoid repeated freeze-thaw cycles.

Reconstitution

It is recommended to reconstitute lyophilized Mouse IL-7-Fc, research grade with deionized sterile-filtered water to a final concentration of 0.1-1.0 mg/mL in a minimal volume of 100 µL. Further dilutions should be prepared with 0.1% bovine serum albumin (BSA) or human serum albumin (HSA) in phosphate-buffered saline.

1.1 Background information

Mouse IL-7-Fc (Fc-fused Mouse IL-7) is a fusion protein consisting of recombinant Mouse IL-7 and the mutated Fc region of human IgG1. The Fc region contains mutations that prevent binding to Fc gamma receptors. Due to the Fc domain forming stable homodimers, the fusion protein presents two IL-7 molecules, one on each Fc arm, resulting in a dimeric configuration of IL-7.

In its function as a stimulatory cytokine, IL-7 is crucial for the development, differentiation and maintenance of T and B cells, and has shown therapeutic potential in boosting immune and antitumor responses. However, IL-7 has a short half-life in vivo and can only act in an autocrine or paracrine manner over short distances. Fusion to the dimeric Fc region extends the protein's half-life in the bloodstream by reducing its clearance rate. While mutations in the Fc region prevent it from binding to Fc gamma receptors (FcγRs), the Fc region can interact with neonatal Fc receptor (FcRn), which protects the protein from degradation and facilitates transcytosis. This results in prolonged circulation time and better tissue distribution, making IL-7-Fc a promising candidate for therapeutic applications.[1,2]

In mice, the administration of Fc-fused IL-7 proteins has been shown to mobilize hematopoietic stem and progenitor cells from the bone marrow,[3] enhance anti-tumor effects in cancer immunotherapy by promoting the expansion of $CD8^+$ cells and have protective effects against viral infections.[4,5] Additionally, the recombinant protein efineptakin alfa (NT-17), another Fc-fused IL-7 variant, has been shown to enhance the expansion, persistence and anti-tumor efficacy of CAR T cells in mice, including in the treatment of solid tumors.[6,7]

Mouse IL-7-Fc can be used for *in vivo* studies in mouse models of disease. The recombinant protein is produced in a human embryonic kidney (HEK) cell line, to ensure proper folding and post-translational modifications.

1.2 Applications

Mouse IL-7-Fc can be used for a variety of applications including:

- Investigating immunomodulatory, antitumor and therapeutic effects in *in vivo* mouse models.
- *In vitro* expansion and priming of T cells.
- *In vitro* differentiation of T cells and iNKT cells.

Optimal concentration for a specific application should be determined by a dose-response experiment.

2. References

1. Ko, S. *et al.* (2025) Engineering FcRn binding kinetics dramatically extends antibody serum half-life and enhances therapeutic potential. J Biol Eng. 19 (1): 35.
2. Kim, A. *et al.* (2024) Understanding the pharmacokinetic journey of Fc-fusion protein, rhIL-7-hyFc using complementary approach of two analytical methods, accelerator mass spectrometry and ELISA. Antib Ther. 7 (2): 105–113.
3. Kim, S. *et al.* (2021) Fc-fused IL-7 mobilizes long-term HSCs in a pro-B cell-dependent manner and synergizes with G-CSF and AMD3100. Leukemia 35 (10): 3030–3034.
4. Kang, M.C. *et al.* (2015) Intranasal Introduction of Fc-Fused Interleukin-7 Provides Long-Lasting Prophylaxis against Lethal Influenza Virus Infection. J Virol. 90 (5): 2273–2284.
5. Kwon, D. *et al.* (2024) Fc-fused IL-7 provides broad antiviral effects against

140-007-318.01

Miltenyi Biotec B.V. & Co. KG
Friedrich-Ebert-Straße 68, 51429 Bergisch Gladbach, Germany
Phone +49 2204 8306-0, Fax +49 2204 85197
macsde@miltenyi.com
www.miltenyibiotec.com

respiratory virus infections through IL-17A-producing pulmonary innate-like T cells. Cell Rep Med. 5 (1): 101362.

6. Kim, M.Y. *et al.* (2022) A long-acting interleukin-7, rhIL-7-hyFc, enhances CAR T cell expansion, persistence, and anti-tumor activity. Nat Commun 13 (1): 3296.
7. Li, D. *et al.* (2024) rhIL-7-hyFc, a long-acting interleukin-7, improves efficacy of CAR-T cell therapy in solid tumors. J Immunother Cancer 12 (7): e008989.

Refer to **www.miltenyibiotec.com** for all data sheets and protocols. Miltenyi Biotec provides technical support worldwide. Visit www.miltenyibiotec.com for local Miltenyi Biotec Technical Support contact information.

Legal notices

Limited product warranty

Miltenyi Biotec B.V. & Co. KG and/or its affiliate(s) warrant this product to be free from material defects in workmanship and materials and to conform substantially with Miltenyi Biotec's published specifications for the product at the time of order, under normal use and conditions in accordance with its applicable documentation, for a period beginning on the date of delivery of the product by Miltenyi Biotec or its authorized distributor and ending on the expiration date of the product's applicable shelf life stated on the product label, packaging or documentation (as applicable) or, in the absence thereof, ONE (1) YEAR from date of delivery ("Product Warranty"). Miltenyi Biotec's Product Warranty is provided subject to the warranty terms as set forth in Miltenyi Biotec's General Terms and Conditions for the Sale of Products and Services available on Miltenyi Biotec's website at www.miltenyibiotec.com, as in effect at the time of order ("Product Warranty"). Additional terms may apply. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING IF A PRODUCT IS SUITABLE FOR CUSTOMER'S PARTICULAR PURPOSE AND APPLICATION METHODS.

Technical information

The technical information, data, protocols, and other statements provided by Miltenyi Biotec in this document are based on information, tests, or experience which Miltenyi Biotec believes to be reliable, but the accuracy or completeness of such information is not guaranteed. Such technical information and data are intended for persons with knowledge and technical skills sufficient to assess and apply their own informed judgment to the information. Miltenyi Biotec shall not be liable for any technical or editorial errors or omissions contained herein.

All information and specifications are subject to change without prior notice. Please contact Miltenyi Biotec Technical Support or visit www.miltenyibiotec.com for the most up-to-date information on Miltenyi Biotec products.

Licenses

This product and/or its use may be covered by one or more pending or issued patents and/or may have certain limitations. Certain uses may be excluded by separate terms and conditions. Please contact your local Miltenyi Biotec representative or visit Miltenyi Biotec's website at www.miltenyibiotec.com for more information.

The purchase of this product conveys to the customer the non-transferable right to use the purchased amount of the product in research conducted by the customer (whether the customer is an academic or for-profit entity). This product may not be further sold. Additional terms and conditions (including the terms of a Limited Use Label License) may apply.

CUSTOMER'S USE OF THIS PRODUCT MAY REQUIRE ADDITIONAL LICENSES DEPENDING ON THE SPECIFIC APPLICATION. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING FOR ITSELF WHETHER IT HAS ALL APPROPRIATE LICENSES IN PLACE. Miltenyi Biotec provides no warranty that customer's use of this product does not and will not infringe intellectual property rights owned by a third party. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

Trademarks

The Miltenyi Biotec logo is a registered trademark or trademark of Miltenyi Biotec B.V. & Co. KG and/or its affiliates in various countries worldwide.

Copyright © 2025 Miltenyi Biotec and/or its affiliates. All rights reserved.

140-007-318.01

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for diagnostic or therapeutic use.