



Miltenyi Biotec

Advanced flow cytometry

FASER Kit – PE

Order No. 130-091-764

Index

1. Description
 - 1.1 Background and product applications
 - 1.2 Reagent requirements
2. General protocol for immunofluorescent staining
3. Example of immunofluorescence amplification using the FASER Kit – PE

Warnings

Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, which is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive conditions may develop.

1. Description

This product is for research use only.

Components	2×2 mL FcR Blocking Reagent, 1 mL PE-Activator (Reagent 1), 1 mL PE-Enhancer (Reagent 2).
Size	For 10 ⁹ total cells, up to 100 stainings.
Staining concentration	FcR Blocking Reagent is used at a dilution of 1:5. The PE-Activator (Reagent 1) and the PE-Enhancer (Reagent 2) are used at a dilution of 1:11.
Product format	Reagents are supplied as a solution containing 0.1% gelatine and 0.05% sodium azide.
Storage	Store protected from light at 4–8 °C. Do not freeze. The expiration date is indicated on the vial label.

1.1 Background and product applications

The FASER (Fluorescence Amplification by Sequential Employment of Reagents) Kit – PE (R-phycoerythrin) is developed to increase the fluorescence intensity of cells labeled with a PE-conjugated antibody. The intensity of the PE fluorescence is amplified by sequential addition of the PE-Activator (Reagent 1) and the PE-Enhancer (Reagent 2). For further enhancement of the fluorescence intensity, sequential addition of the PE-Activator (Reagent 1) and the PE-Enhancer (Reagent 2) can be repeated several times.

The FASER Kit – PE does not affect direct staining with other fluorochrome-conjugated antibodies, but should not be combined with indirect immunofluorescent staining using biotin-conjugated antibodies. The Kit is suitable for PE-labeled, fresh or formaldehyde-fixed cells of any type or species in suspension. Analysis is performed by flow cytometry.


Examples of product application

- Amplification of PE fluorescence which is weak, for example
 - due to PE-conjugated antibody staining of antigens which are expressed at low levels,
 - due to immunomagnetic and PE-conjugated antibody labeling of the same antigen epitope,
 - due to PE-conjugated antibody staining with a low affinity antibody.
- Amplification of the fluorescence signal of a PE-labeled cell fraction for a clearer flow cytometric discrimination from the non-labeled cell fraction.
- Amplification of magnetic labeling with Anti-PE MicroBeads (# 130-048-801) by increasing the number of Anti-PE MicroBead binding sites per cell.

Antibody-labeled cell

1st Amplification round

2nd Amplification round



1.2 Reagent requirements

- Buffer: PBS (phosphate buffered saline) pH 7.2, supplemented with 0.5% BSA (bovine serum albumin) and 2 mM EDTA. Keep buffer cold (4–8 °C).

▲ **Note:** EDTA can be replaced by other supplements such as anticoagulant citrate dextrose formula-A (ACD-A) or citrate phosphate dextrose (CPD). BSA can be replaced by other proteins such as serum albumin, serum or fetal calf serum. Buffers or media containing Ca^{2+} or Mg^{2+} are not recommended for use.

2. General protocol for immunofluorescent staining

▲ If formaldehyde fixation of cells is required, cells should be stained and treated with the FASER Kit – PE prior to fixation.

▲ If the FASER Kit – PE is used for amplification of magnetic labeling with Anti-PE MicroBeads, amplify PE staining as described below. For subsequent magnetic labeling, refer to the Anti-PE MicroBead data sheet.

▲ Volumes for fluorescent labeling given below are for up to 10^7 total cells. When working with fewer than 10^7 cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes, accordingly (e.g. for 2×10^7 total cells, use twice the volume of all indicated reagent volumes and total volumes).

1. Label cells with desired PE-conjugated antibody and afterwards wash cells according to the manufacturer's recommendations.

2. Resuspend up to 10^7 cells in 80 μL of buffer.
3. Add 20 μL of **FcR Blocking Reagent**.
4. Add 10 μL of **PE-Activator (Reagent 1)**.
5. Mix well and incubate for 10 minutes in the dark at 4–8 °C.

▲ **Note:** Working on ice requires increased incubation time. Higher temperatures and/or longer incubation times lead to non-specific cell labeling.


6. Wash cells by adding 1–2 mL of buffer per 10^7 cells and centrifuge at $300 \times g$ for 10 minutes. Pipette off supernatant completely.
7. Resuspend cell pellet in 80 μL of buffer.
8. Add 20 μL of **FcR Blocking Reagent**.
9. Add 10 μL of **PE-Enhancer (Reagent 2)**.
10. Mix well and incubate for 10 minutes in the dark at 4–8 °C.

▲ **Note:** Working on ice requires increased incubation time. Higher temperatures and/or longer incubation times lead to non-specific cell labeling.

11. Wash cells by adding 1–2 mL of buffer per 10^7 cells and centrifuge at $300 \times g$ for 10 minutes. Pipette off supernatant completely.
12. (Optional) Repeat steps 2–11, if further amplification of the PE fluorescence is required.
13. Finally resuspend cell pellet in a suitable amount of buffer for analysis by flow cytometry.

3. Example of immunofluorescence amplification using the FASER Kit – PE

Human peripheral blood mononuclear cells (PBMC) were stained with CD19-PE, human (# 130-091-247), and the PE fluorescence was increased by two rounds of amplification using the FASER Kit – PE. Cells were analyzed by flow cytometry. Cell debris and dead cells were excluded from the analysis based on scatter signals and propidium iodide (PI) fluorescence.



Refer to www.miltenyibiotec.com for all data sheets and protocols. Miltenyi Biotec provides technical support worldwide. Visit www.miltenyibiotec.com for local Miltenyi Biotec Technical Support contact information.

Legal notices

Limited product warranty

Miltenyi Biotec B.V. & Co. KG and/or its affiliate(s) warrant this product to be free from material defects in workmanship and materials and to conform substantially with Miltenyi Biotec's published specifications for the product at the time of order, under normal use and conditions in accordance with its applicable documentation, for a period beginning on the date of delivery of the product by Miltenyi Biotec or its authorized distributor and ending on the expiration date of the product's applicable shelf life stated on the product label, packaging or documentation (as applicable) or, in the absence thereof, ONE (1) YEAR from date of delivery ("Product Warranty"). Miltenyi Biotec's Product Warranty is provided subject to the warranty terms as set forth in Miltenyi Biotec's General Terms and Conditions for the Sale of Products and Services available on Miltenyi Biotec's website at www.miltenyibiotec.com, as in effect at the time of order ("Product Warranty"). Additional terms may apply. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING IF A PRODUCT IS SUITABLE FOR CUSTOMER'S PARTICULAR PURPOSE AND APPLICATION METHODS.

Technical information

The technical information, data, protocols, and other statements provided by Miltenyi Biotec in this document are based on information, tests, or experience which Miltenyi Biotec believes to be reliable, but the accuracy or completeness of such information is not guaranteed. Such technical information and data are intended for persons with knowledge and technical skills sufficient to assess and apply their own informed judgment to the information. Miltenyi Biotec shall not be liable for any technical or editorial errors or omissions contained herein.

All information and specifications are subject to change without prior notice. Please contact Miltenyi Biotec Technical Support or visit www.miltenyibiotec.com for the most up-to-date information on Miltenyi Biotec products.

Licenses

This product and/or its use may be covered by one or more pending or issued patents and/or may have certain limitations. Certain uses may be excluded by separate terms and conditions. Please contact your local Miltenyi Biotec representative or visit Miltenyi Biotec's website at www.miltenyibiotec.com for more information.

The purchase of this product conveys to the customer the non-transferable right to use the purchased amount of the product in research conducted by the customer (whether the customer is an academic or for-profit entity). This product may not be further sold. Additional terms and conditions (including the terms of a Limited Use Label License) may apply.

CUSTOMER'S USE OF THIS PRODUCT MAY REQUIRE ADDITIONAL LICENSES DEPENDING ON THE SPECIFIC APPLICATION. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING FOR ITSELF WHETHER IT HAS ALL APPROPRIATE LICENSES IN PLACE. Miltenyi Biotec provides no warranty that customer's use of this product does not and will not infringe intellectual property rights owned by a third party. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

Trademarks

MACS and the Miltenyi logo are registered trademarks or trademarks of Miltenyi Biotec and/or its affiliates in various countries worldwide.

Copyright © 2021 Miltenyi Biotec and/or its affiliates. All rights reserved.