Colon TumorMACS™ Medium
human

Order no. 130-127-169

Contents

1. Description
   - 1.1 Background information
   - 1.2 Applications
2. Protocol
   - 2.1 Preparation of complete medium
   - 2.2 Culture of tumor cells

1. Description

This product is for research use only.

Components	**500 mL TumorMACS Basal Medium, human** **10 mL Colon TumorMACS Supplement (50×), human**
Specifications	Osmolality: 300–340 mOsmol/kg pH: 7.2–7.4
Storage	Store TumorMACS Basal Medium, human protected from light at +2 to +8 °C. Do not freeze. The expiration date is indicated on the bottle label. Upon arrival store Colon TumorMACS Supplement (50×), human at –20 °C. Aliquots of the supplement should be stored at –20 °C. Avoid repeated freeze-thaw cycles. The expiration date is indicated on the vial label.

1.1 Background information

TumorMACS Media are optimized, standardized, and serum-free media for the reliable expansion of tumor cells from primary or xenotransplanted human tumor samples. Taking into account that different tumor entities have specific culture requirements, Miltenyi Biotec has developed tumor-specific TumorMACS Supplements. Accordingly, the appropriate specific medium should be used to achieve optimal cell culture results for a particular tumor entity. Tumor cells from freshly dissociated tissue can be cultured over multiple passages, thus allowing for *in vitro* assays, such as drug screenings on primary cell cultures or characterization studies including gene expression or protein profiling. In addition, TumorMACS Media support the establishment of new cell lines and the generation of cell numbers that are sufficiently large for further experiments, such as *in vivo* transplantation studies in animals. TumorMACS Media efficiently expand cultures of primary tumor cells *in vitro* while maintaining the cell populations' heterogeneity and tumorigenic potential. The media are manufactured under controlled conditions using ingredients of high quality. TumorMACS Media offer consistent lot-to-lot performance and optimal conditions for the cultivation of primary tumor cells.

1.2 Applications

- Generation and expansion of tumor cell cultures from primary or xenografted human colorectal tumors.
- Derivation of primary tumor cell lines from primary or xenografted human tumors.
- The medium supports the transition of tumor cells into physiologically relevant 3D culture formats, including tumor organoids and tumoroids. For more information, please refer to www.miltenyibiotec.com/tumoroids.
- A detailed guideline for the initiation of primary tumor cell cultures is available at www.miltenyibiotec.com/tumorculture.

2. Protocol

2.1 Preparation of complete medium

Before Colon TumorMACS Medium can be used in cell culture, the two media components need to be mixed according to the following protocol to obtain the complete medium.

1. Thaw Colon TumorMACS Supplement (50×) at room temperature prior to use. Thawed material must be used immediately or aliquoted to the preferred sample volume, for example, 1 mL. Store aliquots at –20 °C. Avoid additional freeze-thaw cycles.
2. To obtain the complete medium add 10 mL Colon TumorMACS Supplement (50×) to 500 mL TumorMACS Basal Medium. The medium is ready-to-use now. Use the complete medium within 2 weeks when stored at +2 to +8 °C.

2.2 Culture of tumor cells

1. Plate 2×10^5 cells/cm² in TumorMACS Medium.
2. Incubate at +37 °C and 5–7.5% CO_2.
3. Change medium every 4–5 days or as soon as it turns yellowish.
4. Split or remove fibroblasts at a confluency of 80–90%.

Refer to **www.miltenyibiotec.com** for all data sheets and protocols. Miltenyi Biotec provides technical support worldwide. Visit www.miltenyibiotec.com for local Miltenyi Biotec Technical Support contact information.

Miltenyi Biotec B.V. & Co. KG
Friedrich-Ebert-Straße 68, 51429 Bergisch Gladbach, Germany
Phone +49 2204 8306-0, Fax +49 2204 85197
macsde@miltenyi.com
www.miltenyibiotec.com

140-005-982.04

Legal notices

Limited product warranty

Miltenyi Biotec B.V. & Co. KG and/or its affiliate(s) warrant this product to be free from material defects in workmanship and materials and to conform substantially with Miltenyi Biotec's published specifications for the product at the time of order, under normal use and conditions in accordance with its applicable documentation, for a period beginning on the date of delivery of the product by Miltenyi Biotec or its authorized distributor and ending on the expiration date of the product's applicable shelf life stated on the product label, packaging or documentation (as applicable) or, in the absence thereof, ONE (1) YEAR from date of delivery ("Product Warranty"). Miltenyi Biotec's Product Warranty is provided subject to the warranty terms as set forth in Miltenyi Biotec's General Terms and Conditions for the Sale of Products and Services available on Miltenyi Biotec's website at www.miltenyibiotec.com, as in effect at the time of order ("Product Warranty"). Additional terms may apply. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING IF A PRODUCT IS SUITABLE FOR CUSTOMER'S PARTICULAR PURPOSE AND APPLICATION METHODS.

Technical information

The technical information, data, protocols, and other statements provided by Miltenyi Biotec in this document are based on information, tests, or experience which Miltenyi Biotec believes to be reliable, but the accuracy or completeness of such information is not guaranteed. Such technical information and data are intended for persons with knowledge and technical skills sufficient to assess and apply their own informed judgment to the information. Miltenyi Biotec shall not be liable for any technical or editorial errors or omissions contained herein.

All information and specifications are subject to change without prior notice. Please contact Miltenyi Biotec Technical Support or visit www.miltenyibiotec.com for the most up-to-date information on Miltenyi Biotec products.

Licenses

This product and/or its use may be covered by one or more pending or issued patents and/or may have certain limitations. Certain uses may be excluded by separate terms and conditions. Please contact your local Miltenyi Biotec representative or visit Miltenyi Biotec's website at www.miltenyibiotec.com for more information.

The purchase of this product conveys to the customer the non-transferable right to use the purchased amount of the product in research conducted by the customer (whether the customer is an academic or for-profit entity). This product may not be further sold. Additional terms and conditions (including the terms of a Limited Use Label License) may apply.

CUSTOMER'S USE OF THIS PRODUCT MAY REQUIRE ADDITIONAL LICENSES DEPENDING ON THE SPECIFIC APPLICATION. THE CUSTOMER IS SOLELY RESPONSIBLE FOR DETERMINING FOR ITSELF WHETHER IT HAS ALL APPROPRIATE LICENSES IN PLACE. Miltenyi Biotec provides no warranty that customer's use of this product does not and will not infringe intellectual property rights owned by a third party. BY USE OF THIS PRODUCT, THE CUSTOMER AGREES TO BE BOUND BY THESE TERMS.

Trademarks

The Miltenyi Biotec logo and TumorMACS are registered trademarks or trademarks of Miltenyi Biotec B.V. & Co. KG and/or its affiliates in various countries worldwide. All other trademarks mentioned in this publication are the property of their respective owners and are used for identification purposes only.

Copyright © 2026 Miltenyi Biotec and/or its affiliates. All rights reserved.